

RECOGNIZING HYPERAMMONEMIA



CHECK AMMONIA

HYPERAMMONEMIA CAN BE EXTREMELY TOXIC TO THE CENTRAL NERVOUS SYSTEM IN INFANTS, CHILDREN, AND ADULTS.

HELP PROTECT THE BRAIN

HYPERAMMONEMIA OVERVIEW

Hyperammonemia—a metabolic condition characterized by excess ammonia in the blood—can affect patients at any age.¹

In the neonate (days 1-28), hyperammonemia is a rare, life-threatening problem that requires prompt intervention.²

The clinical presentation of neonatal hyperammonemia, which can mimic sepsis, includes non-specific symptoms that are mainly neurological in origin.^{1,3}

Hyperammonemia in infants, children, and adults may be more episodic, difficult to recognize, and precipitated by catabolic events, protein overload, or certain drugs.^{4,5,6}

Although their hyperammonemia may be less severe, patients may nonetheless face serious health risks.^{4,7} Nonspecific symptoms may be attributed to a wide variety of neurological and psychiatric disorders.^{8,9}

The causes of hyperammonemia are diverse. Some of the more common causes include liver disease, reactions to drugs, hemolytic disease, gastrointestinal bleeds, and urea cycle disorders (UCDs) or other inborn errors of metabolism (IEMs).^{3,10,11}

Untreated hyperammonemia can result in irreversible brain damage and death.¹ To improve the prognosis of patients, an increased awareness of hyperammonemia is needed.¹ Knowledge of common symptoms and triggers, along with prompt ammonia testing and treatment, is critical for improved outcomes.^{1,5}

GO BEYOND SEPSIS: THINK INBORN ERRORS OF METABOLISM

Symptoms of sepsis mimic those of hyperammonemia.³ However, when severe symptoms develop after 24 hours of age, the cause is usually a UCD or other IEM.¹¹

Do not wait for sepsis results.¹

In neonates with any unexplained alteration in consciousness or encephalopathy¹:

- Check ammonia levels
- Call a metabolic geneticist

CHECK AMMONIA

HYPERAMMONEMIA CAN BE EXTREMELY TOXIC
TO THE CENTRAL NERVOUS SYSTEM IN INFANTS, CHILDREN, AND ADULTS.

HELP PROTECT THE BRAIN

HYPERAMMONEMIA IN NEONATES

The clinical presentation of neonatal hyperammonemia may include many symptoms that are neurological in origin.¹

Signs and symptoms may include:

Neurological	Gastrointestinal*
Altered level of consciousness (from somnolence to lethargy to coma) —mimicking encephalitis or drug intoxication ⁶	Progressive poor appetite ⁶ or poor feeding ^{1,13}
Irritability ¹²	Vomiting ⁶
Acute encephalopathy, ⁶ cerebral edema ¹²	
Seizures (generally not isolated but with altered level of consciousness) ^{6,13}	
Hypotonia ¹⁴	
Neurologic posturing ¹⁵	
Hepatomegaly ¹²	
Multiorgan failure ⁶	
Peripheral circulatory failure ⁶	
Respiratory distress, ⁶ tachypnea, ¹² hyperventilation, ⁶ respiratory alkalosis ¹	
Sepsis-like picture ^{6,14} (including temperature instability, ⁶ hypo- or hyperthermia ¹)	

*Gastrointestinal symptoms are neurological in origin.¹

CHECK AMMONIA IMMEDIATELY IN NEONATES WITH UNEXPLAINED^{1,6*}

- Poor feeding or vomiting
- Alteration in consciousness
- Encephalopathy
- Respiratory distress
- Sepsis-like symptoms

HYPERAMMONEMIA IN INFANTS, CHILDREN, AND ADULTS

The clinical presentation of hyperammonemia beyond the neonatal period may be less severe but more variable and episodic.^{4,6} Because symptoms are non-specific, they may be attributed to a range of psychiatric or neurological disorders.^{8,9} These symptoms may instead indicate a late-onset urea cycle disorder (UCD).

Signs and symptoms may include:

Neurological	Gastrointestinal	Psychiatric
Confusion, lethargy, dizziness ⁶	Abdominal pain, ⁶ nausea, ^{13,14} vomiting ¹	Behavioral changes, mood alteration, hyperactivity, aggressiveness, ⁶ combativeness ¹³
Migraine-like headaches ⁶	Protein aversion, self-selected low-protein diet ⁶	Delusions, psychosis ⁴
Tremor, ataxia, dysarthria ⁶	Failure to thrive ⁶	Sleep disorders ¹⁵
Intellectual/ learning disabilities, neurodevelopmental delay ⁶	Hepatomegaly, elevated liver enzymes ⁶	
Seizures ¹		
Hemiplegia ³		
Coma ¹⁶		

CHECK AMMONIA IMMEDIATELY IN INFANTS, CHILDREN, AND ADULTS WITH UNEXPLAINED^{1,6,7,12,17}:


- GI presentations (e.g., vomiting, protein aversion)
- Alteration in consciousness
- Encephalopathy
- Movement disorders or seizures
- Learning problems or developmental delay
- Psychiatric presentations

LATE-ONSET UCDs AND HYPERAMMONEMIA

Hyperammonemia due to a late-onset urea cycle disorder (UCD) can occur at any age.^{1,5} Adult patients, in particular, are being diagnosed with UCDs with greater frequency.⁵

Clinical Presentation of UCDs*

Most patients with UCDs present with symptoms outside the newborn period.⁷



*Data from a longitudinal study of 614 UCD patients conducted by the urea cycle disorders consortium (UCDC), a member of the NIH Rare Diseases Clinical Research Network.⁷

Patients with milder UCDs may function relatively normally—for months or years—before their UCD is clinically recognized, although they may have experienced multiple episodes of mild to moderately severe hyperammonemia.^{5,18,19} These patients typically face serious health risks, including a significant risk for developmental disabilities and a 10% risk of mortality.⁷

GO BEYOND LIVER DISEASE: THINK INBORN ERRORS OF METABOLISM

In patients with any unexplained alteration in consciousness or encephalopathy, including patients with hepatic disease^{1,20}:

- Consider late-onset UCDs or other IEMs
- Check ammonia levels
- Call a metabolic geneticist

**FAILURE TO RECOGNIZE HYPERAMMONEMIA
CAN LEAD TO BRAIN DAMAGE OR DEATH.⁴**



HYPERAMMONEMIA TRIGGERS

A precipitating catabolic event can trigger a hyperammonemic crisis in patients with a urea cycle disorder (UCD) or other underlying metabolic disorder.^{4,5}

Known triggers may include:

- Traumatic injury or surgery^{4,20}
- Illness, including viral or bacterial infections, fever, or vomiting^{1,6}
- Internal bleeding⁶
- Protein overload, such as eating barbecue⁶
- Decreased protein intake, such as fasting pre-surgery⁶
- Medications
 - Valproate⁶
 - Certain chemotherapy drugs⁶
 - High-dose glucocorticoids⁶
 - Salicylic acid or aspirin (aspirin use in children with viral illness may cause Reye syndrome)^{1,21}
- Rapid growth²²
- Prolonged or intense physical exercise, such as bodybuilding^{6,23}
- Peripartum stressors¹⁵

REFERENCES

1. Haberle J. Clinical practice: The management of hyperammonemia. *Eur J Pediatr* 2011; 170:21-34.
2. Vergano SA, Crossette JM, Cusick FC, Desai BR, Deardorff MA, Sondheimer N. Improving surveillance for hyperammonemia in the newborn. *Mol Genet Metab* 2013;110: 102-105.
3. Broomfield A, Grunewald S. How to use serum ammonia. *Arch Dis Child Educ Pract Ed* 2012;97:72-77.
4. Summar ML, Tuchman M. Proceedings of a consensus conference for the management of patients with urea cycle disorders. *J Pediatr* 2001;138 (1 Suppl):S6-S10.
5. Summar ML, Barr F, Dawling S, Smith W, Lee B, Singh RH, Rhead WJ, Sniderman King L, Christman BW. Unmasked adult-onset urea cycle disorders in the critical care setting. *Crit Care Clin* 2005;21: S1-S8.
6. Haberle J, Boddaert N, Burlina A, Chakrapani A, Dixon M, Huemer M, Karall D, Martinelli D, Sanjurjo Crespo P, Santer R, Servais A, Valayannopoulos V, Lindner M, Rubio V, Dionisi-Vici C. Suggested guidelines for the diagnosis and management of urea cycle disorders. *Orphanet J Rare Dis* 2012;7:32.
7. Batshaw ML, Tuchman M, Summar M, Seminara J. A longitudinal study of urea cycle disorders. *Mol Genet Metab* 2014;113: 127-130.
8. Gascon-Bayarri J, Campdelacreu J, Estela J, Rene R. Severe hyperammonemia in late-onset ornithine transcarbamylase deficiency triggered by steroid administration. *Case Rep Neurol Med* 2015;1-3.
9. Serrano M, Martins C, Perez-Duenas B, Gomez-Lopez L, Murgui E, Fons C, Garcia-Cazorla A, Artuch R, Jara F, Arranz J, Haberle J, Briones P, Campistol J, Pineda M, Vilaseca M. Neuropsychiatric manifestations in late-onset urea cycle disorder patients. *J Child Neurol* 2010;25(3):352-358.
10. Orton DJ, Gifford JL, Seiden-Long I, Khan A, de Koning L. Critically high plasma ammonia in an adolescent girl. *Clin Chem* 2016;62(12):1565-1569.
11. Burton BK. Inborn errors of metabolism in infancy: A guide to diagnosis. *Pediatrics* 1998;102(6):E69.
12. Al Kaabi EH, El-Hattab AW. N-acetylglutamate synthase deficiency: Novel mutation associated with neonatal presentation and literature review of molecular and phenotypic spectra. *Mol Genet Metab Rep* 2016;8:94-98.
13. Ah Mew N, Caldovic L. N-acetylglutamate synthase deficiency: An insight into the genetics, epidemiology, pathophysiology, and treatment. *Appl Clin Gen* 2011;4:127-135.
14. Cartagena A, Prasad AN, Rupar CA, Strong M, Tuchman M, Ah Mew N, Prasad C. Recurrent encephalopathy: NAGS (N-acetylglutamate synthase) deficiency in adults. *Can J Neurol Sci* 2013;40:3-9.
15. Ah Mew N, Lanpher BC, Gropman A, Chapman KA, Simpson KL, Urea Cycle Disorders Consortium, Summar ML. Urea cycle disorders overview. In Pagon RA, Adam MP, Ardinger HH et al, eds. *GeneReviews@ [Internet]*. Seattle, WA: University of Washington, Seattle; 1993-2017. <http://www.ncbi.nlm.nih.gov/books/NBK1217>. Updated June 22, 2017. Accessed September 22, 2017.
16. Caldovic L, Morizono H, Tuchman M. Mutations and polymorphisms in the human N-acetylglutamate synthase (NAGS) gene. *Hum Mutat* 2007;28:754-759.
17. Kolker S, Cazorla AG, Valayannopoulos V, Lund AM, Burlina AB, Sykut-Cegielska J, Wijburg W, Teles EL, Zeman J, Dionisi-Vici C, Baric I, Karall D, Augoustides-Savvopoulou P, Aksglaede L et al. The phenotypic spectrum of organic acidemias and urea cycle disorders. Part 1: the initial presentation. *J Inheret Metab Dis* 2015;38:1041-1057.
18. Summar ML, Dobbelaere D, Brusilow S, Lee B. Diagnosis, symptoms, frequency and mortality of 260 patients with urea cycle disorders from a 21-year, multicenter study of acute hyperammonaemic episodes. *Acta Pediatr* 2008;97(10):1420-1425.
19. Summar ML. *The Physician's Guide to Urea Cycle Disorders*. The National Organization for Rare Disorders. NORD Guides for Physicians series. 2012. Available at: <https://rarediseases.org/physician-guide/urea-cycle-disorders/>. Accessed September 21, 2017.
20. LaBuzetta JN, Yao JZ, Bourque DL, Zivin J. Adult nonhepatic hyperammonemia: a case report and differential diagnosis. *Am J Med* 2010;123(10):885-891.
21. N-acetylglutamate synthetase deficiency. The National Organization for Rare Disorders. 2016. Available at: <https://rarediseases.org/rare-diseases/n-acetylglutamate-synthetase-deficiency>. Accessed September 11, 2017.
22. Smith W, Kishnani PS, Lee B, Singh RH, Rhead WJ, Sniderman King L, Smith M, Summar M. Urea cycle disorders: clinical presentation outside the newborn period. *Crit Care Clin* 2005;21: S9-S17.
23. Panlaqui OM, Tran K, Johns A, McGill J, White H. Acute hyperammonemic encephalopathy in adult onset ornithine transcarbamylase deficiency. *Intensive Care Med* 2008;34:1922-1924.
24. Auron A, Brophy PD. Hyperammonemia in review: pathophysiology, diagnosis, and treatment. *Pediatr Nephrol* 2012;27: 207-222.

HYPERAMMONEMIA: A MEDICAL EMERGENCY

The challenge is early detection.

Delayed diagnosis or treatment of hyperammonemia, regardless of cause, can lead to neurological damage and potentially a fatal outcome.²⁴

Hyperammonemia in neonates:

- Initial non-specific symptoms often mimic sepsis.³
- Newborns who develop severe hyperammonemia >24 hours of age usually have a urea cycle disorder (UCD) or other inborn error of metabolism.¹¹

Hyperammonemia in infants, children, and adults:

- More than two-thirds of UCD patients present later in life.⁷
- Catabolic events, protein overload, or certain drugs may cause a hyperammonemic crisis in patients with an underlying UCD.⁶

ACT QUICKLY TO IDENTIFY HYPERAMMONEMIA:

- Check ammonia levels
- Call a metabolic geneticist

CHECK AMMONIA

HYPERAMMONEMIA CAN BE EXTREMELY TOXIC
TO THE CENTRAL NERVOUS SYSTEM IN INFANTS, CHILDREN, AND ADULTS.

HELP PROTECT THE BRAIN

VISIT [CHECKAMMONIA.COM](https://checkammonia.com)



National Urea Cycle Disorders Foundation
Families • Research • Education • Support ... Hope